

A general theory of self-similarity II: recognition

Tom Leinster

Department of Mathematics, University of Glasgow
www.maths.gla.ac.uk/~tl
tl@maths.gla.ac.uk

Abstract

This paper concerns the self-similarity of topological spaces, in the sense defined in [Lei1]. I show how to recognize self-similar spaces, or more precisely, universal solutions of self-similarity systems. Examples include the standard simplices (self-similar by barycentric subdivision) and solutions of iterated function systems. Perhaps surprisingly, every compact metrizable space is self-similar in at least one way. From this follow the classical results on the role of the Cantor set among compact metrizable spaces.

Contents

Introduction	2
1 Recognition theorems	2
2 Examples	8
3 Spaces admitting a self-similarity	17
4 The Cantor set	21
References	28

Introduction

This paper's predecessor [Lei1] introduced the notions of 'self-similarity system' (to be thought of as a system of simultaneous equations in which each member of a family of objects is expressed as a gluing-together of other members) and 'universal solution' of such a system. A self-similarity system possesses a universal solution if and only if an explicit condition **S** holds; if so then the universal solution is unique and can be constructed explicitly.

Few examples have been given so far. In principle one can take any self-similarity system $(\mathbb{A}, M)$ satisfying **S** and find the universal solution by going through the explicit construction. In practice this is cumbersome and it is much quicker to apply one of the Recognition Theorems proved in §1 below, as follows.

Recall that if (J, γ) is a universal solution of $(\mathbb{A}, M)$ then $J \cong M \otimes J$, that is, each space Ja must be homeomorphic to the prescribed gluing-together of (Jb) s. This necessary condition is not sufficient (see [Lei1], after Definition 1.9), but it turns out to become sufficient if, roughly speaking, each space Ja is nonempty, compact, and made up of *smaller* copies of (Jb) s. Put another way, a solution (J, γ) of the system is universal if it consists of nonempty compact metric spaces and contractions between them. This is the Crude Recognition Theorem, essentially; it gives sufficient conditions for an M -coalgebra to be a universal solution. A more refined result is the Precise Recognition Theorem, which gives necessary and sufficient conditions.

To find the universal solution of a given self-similarity system, one can therefore make a guess and check that it satisfies the hypotheses of a Recognition Theorem. I do this in §2 for various specific self-similarity systems, formulating, for instance, the self-similarity of the cubes $[0, 1]^n$, the simplices Δ^n (by both barycentric and edgewise subdivision), the Cantor set, and the solutions of certain iterated function systems.

A further challenge is to recognize which topological spaces arise as universal solutions to some self-similarity system. The Recognition Theorems show that all such spaces are compact and metrizable, so the question is: which compact metrizable spaces are self-similar? Perhaps surprisingly, the answer turns out to be: all of them (§3).

A measure of the non-triviality of this result is that some of the classical results on the Cantor set can be derived as corollaries: for instance, that every nonempty compact metric space is a quotient of the Cantor set. This is the subject of §4.

Terminology and notation from [Lei1] are used without further mention. An overview of [Lei1] and the present paper is available: [Lei2].

I gratefully acknowledge a Nuffield Foundation award NUF-NAL 04.

1 Recognition theorems

Fix a self-similarity system $(\mathbb{A}, M)$.

If $(\mathbb{A}, M)$ has a universal solution then by Theorem A.1 of [Lei1], condition

S holds and the universal solution is the (I, ι) constructed in §4 of [Lei1]. We begin by establishing various properties of (I, ι) , working throughout in **Top** rather than **Set**.

First, Lambek's Lemma guarantees that (I, ι) is a **fixed point** of M , that is, an M -coalgebra whose structure map is an isomorphism. (Note that by definition, fixed points are nondegenerate.) A fixed point (J, γ) is a coalgebra, but can also be regarded as an algebra (J, ψ) where $\psi = \gamma^{-1}$. By definition, an M -**algebra** (in **Top**) is a nondegenerate functor $J : \mathbb{A} \longrightarrow \mathbf{Top}$ together with a map $\psi : M \otimes J \longrightarrow J$; but

$$(M \otimes J)a = \int^b M(b, a) \times Jb,$$

so ψ consists of a map $\psi_m : Jb \longrightarrow Ja$ for each module element $m : b \dashrightarrow a$, satisfying naturality axioms. These say that if

$$b' \xrightarrow{g} b \dashrightarrow^m a \xrightarrow{f} a'$$

then $\psi_{fmg} = (Jf) \circ \psi_m \circ (Jg)$.

For example, the fixed point (I, ι) has algebra structure $\phi = \iota^{-1}$, where the components ϕ_m are as defined in [Lei1] (after Lemma 6.2).

Lemma 1.1 *Let (J, ψ) be a fixed point of M . Then for each module element $b \dashrightarrow^m a$, the map $Jb \xrightarrow{\psi_m} Ja$ is closed.*

Proof ψ_m is the composite

$$Jb \xrightarrow{m \otimes -} (M \otimes J)a \xrightarrow[\sim]{\psi_a} Ja$$

and $m \otimes -$ is closed by [Lei1, 6.14] and nondegeneracy of J . $\square$

Second, the sets Ia are not empty unless they are forced to be. If, for instance, $a \in \mathbb{A}$ and there is no module element of the form $b \dashrightarrow^m a$, then for any coalgebra (X, ξ) we have $\xi_a : Xa \longrightarrow (M \otimes X)a = \emptyset$, so Xa is forced to be empty. More generally, Xa is forced to be empty if there is no infinite chain $\cdots \dashrightarrow a_1 \dashrightarrow a$. Say that a functor $X : \mathbb{A} \longrightarrow \mathbf{Set}$ is **occupied** if $Ja \neq \emptyset \Rightarrow Xa \neq \emptyset$ for all $a \in \mathbb{A}$. Trivially, I is occupied.

Third, the spaces Ia are metrizable, by

Lemma 1.2 *A compact space is metrizable if and only if it is Hausdorff and second countable.*

Proof See [Bou2, IX.2.9] (where ‘compact’ means compact Hausdorff). $\square$

One naturally asks how a metric can be defined. There are many possible metrics and apparently no canonical choice among them, but the following result tells us all we need to know. Recall from [Lei1] (before Lemma 6.3) that for each $a \in \mathbb{A}$ and $n \in \mathbb{N}$ we have a closed binary relation R_n^a on Ia .

Proposition 1.3 (Metric on Ia) *For each $a \in \mathbb{A}$ there exists a metric d on Ia compatible with the topology on Ia and such that*

$$\forall \varepsilon > 0, \exists n \in \mathbb{N} : (t, t') \in R_n^a \Rightarrow d(t, t') \leq \varepsilon.$$

Proof In fact, the stated property holds for any compatible metric d on Ia (and by Lemma 1.2, there is at least one such). For if it does not hold, there is some $\varepsilon > 0$ such that $d^{-1}[\varepsilon, \infty) \cap R_n^a \neq \emptyset$ for all $n \in \mathbb{N}$, where $d^{-1}[\varepsilon, \infty)$ is the inverse image of $[\varepsilon, \infty)$ under $d : Ia \times Ia \longrightarrow \mathbb{R}$. Since d is continuous, $d^{-1}[\varepsilon, \infty)$ is closed. Each R_n^a is also closed and Ia is compact, so $d^{-1}[\varepsilon, \infty) \cap \bigcap_{n \in \mathbb{N}} R_n^a \neq \emptyset$; but by Lemma 6.3 of [Lei1], $\bigcap_{n \in \mathbb{N}} R_n^a = \Delta_{Ia}$, a contradiction. $\square$

Theorem 1.4 (Precise Recognition Theorem) *The following are equivalent conditions on a fixed point $(J, \gamma = \psi^{-1})$ of M in **Top**:*

- a. (J, γ) is a universal solution of $(\mathbb{A}, M)$ in **Top**
- b. J is occupied, and for each $a \in \mathbb{A}$ the space Ja is compact and can be metrized in such a way that

$$\forall \varepsilon > 0, \exists n \in \mathbb{N} : \forall (a_n \xrightarrow{m_n} \cdots \xrightarrow{m_1} a_0) \in \mathcal{I}_n a, \\ \text{diam}(\psi_{m_1} \cdots \psi_{m_n}(Ja_n)) \leq \varepsilon$$

- c. J is occupied, and for each $a \in \mathbb{A}$, the space Ja is compact and

$$\forall (\cdots \xrightarrow{m_1} a_0) \in \mathcal{I}a, \left| \bigcap_{n \in \mathbb{N}} \psi_{m_1} \cdots \psi_{m_n}(Ja_n) \right| \leq 1$$

(where $| \cdot |$ means cardinality).

Proof

(a) $\Rightarrow$ (b) Assume (a). By Theorem A.1 of [Lei1], condition **S** holds, so (J, γ) is the universal solution (I, ι) constructed in §4 of [Lei1]. Certainly I is occupied and each Ia is compact. Proposition 1.3 then gives metrics with exactly the property required.

(b) $\Rightarrow$ (c) Trivial.

(c) $\Rightarrow$ (a) Assume (c). First I claim that each Ja is Hausdorff. For each $n \in \mathbb{N}$, put

$$\tilde{R}_n^a = \bigcup \{ (\psi_{m_1} \cdots \psi_{m_n}(Ja_n))^2 \mid (a_n \xrightarrow{m_n} \cdots \xrightarrow{m_1} a_0) \in \mathcal{I}_n a \} \subseteq Ja \times Ja.$$

Then each $\tilde{R}_n^a$ is closed by Lemma 1.1 and finiteness of $\mathcal{J}_n a$, so it suffices to show that $\bigcap_{n \in \mathbb{N}} \tilde{R}_n^a = \Delta_{Ja}$. Certainly $\bigcap_{n \in \mathbb{N}} \tilde{R}_n^a \supseteq \Delta_{Ja}$. Conversely, let $(t, t') \in \bigcap_{n \in \mathbb{N}} \tilde{R}_n^a$. For each $n \in \mathbb{N}$ we may choose

$$(a_n^n \xrightarrow{m_n^n} \cdots \xrightarrow{m_1^n} a_0^n) \in \mathcal{J}_n a$$

such that $t, t' \in \psi_{m_1^n} \cdots \psi_{m_n^n}(Ja_n^n)$. By König's Lemma [Lei1, 5.1] and finiteness of $\mathcal{J}_n a$, we may then choose

$$(\cdots \xrightarrow{m_2} a_1 \xrightarrow{m_1} a_0) \in \mathcal{J} a$$

such that for all $r \in \mathbb{N}$, there exists $n \geq r$ satisfying

$$(a_r \xrightarrow{m_r} \cdots \xrightarrow{m_1} a_0) = (a_r^n \xrightarrow{m_r^n} \cdots \xrightarrow{m_1^n} a_0^n).$$

Hence

$$t, t' \in \bigcap_{r \in \mathbb{N}} \psi_{m_1} \cdots \psi_{m_r}(Ja_r),$$

which by (c) implies that $t = t'$, as required.

The next part of the strategy is to show that (J, γ) is the universal solution in **Set**. Let (X, ξ) be an M -coalgebra in **Set**. Define for each $a \in \mathbb{A}$ and $x \in Xa$ a sequence $(K_n(x))_{n \in \mathbb{N}}$ of subsets of Ja as follows:

- $K_0(x) = Ja$
- $K_{n+1}(x) = \bigcap \{ \psi_m K_n(y) \mid b \xrightarrow{m} a, y \in Xb, \xi(x) = m \otimes y \}.$

We eventually show that $\bigcap_{n \in \mathbb{N}} K_n(x)$ is a one-element set and that its element is the value at x of the unique map $(X, \xi) \longrightarrow (J, \gamma)$.

Claim $K_n(x) \supseteq K_{n+1}(x)$ for each $a \in \mathbb{A}$, $x \in Xa$, $n \in \mathbb{N}$.

Proof For $n = 0$ this is trivial. For $n \geq 0$, inductively,

$$\begin{aligned} K_{n+1}(x) &= \bigcap \{ \psi_m K_n(y) \mid \xi(x) = m \otimes y \} \\ &\supseteq \bigcap \{ \psi_m K_{n+1}(y) \mid \xi(x) = m \otimes y \} \\ &= K_{n+2}(x). \end{aligned}$$

Claim $(Jf)K_n(x) \subseteq K_n(fx)$ for each $(a \xrightarrow{f} a')$ in $\mathbb{A}$, $x \in Xa$, $n \in \mathbb{N}$.

Proof For $n = 0$ this is trivial. Suppose inductively that the claim holds for $n \in \mathbb{N}$. Suppose that $t \in K_{n+1}(x)$ and that $b' \xrightarrow{m'} a'$ and $y' \in Xb$ with $\xi(fx) = m' \otimes y'$; we have to show that $ft \in \psi_{m'} K_n(y')$.

We may choose $b \xrightarrow{m} a$ and $y \in Xb$ such that $\xi(x) = m \otimes y$. Then $m' \otimes y' = \xi(fx) = fm \otimes y$, so there exist a commutative square

$$\begin{array}{ccc} & c & \\ g \swarrow & & \searrow g' \\ b & & b' \\ fm \searrow & & \swarrow m' \\ & a' & \end{array}$$

and $z \in Xc$ such that $y = gz$ and $y' = g'z$. Now $\xi(x) = m \otimes y = m \otimes gz = mg \otimes z$, so $t \in \psi_{mg} K_n(z)$, so

$$\begin{aligned} ft &\in \psi_{fmg} K_n(z) = \psi_{m'g'} K_n(z) = \psi_{m'}(Jg') K_n(z) \\ &\subseteq \psi_{m'} K_n(g'z) = \psi_{m'} K_n(y') \end{aligned}$$

(the penultimate step by inductive hypothesis), as required.

Claim $K_n(x)$ is closed in Ja for each $a \in \mathbb{A}$, $x \in Xa$, $n \in \mathbb{N}$.

Proof Lemma 1.1 and induction.

Claim $K_n(x)$ is nonempty for each $a \in \mathbb{A}$, $x \in Xa$, $n \in \mathbb{N}$.

Proof We use induction on $n \in \mathbb{N}$ for all $a \in \mathbb{A}$ and $x \in Xa$ simultaneously.

For $n = 0$, let $a \in \mathbb{A}$ and $x \in Xa$. There exists a resolution of x , and in particular an element of Ja . Since J is occupied, $\emptyset \neq Ja = K_0(x)$.

Now let $n \in \mathbb{N}$, $a \in \mathbb{A}$, and $x \in Xa$; we have to prove that $K_{n+1}(x)$ is nonempty. Since $K_{n+1}(x)$ is an intersection of a family of closed subsets of a compact space, it suffices to show that the intersection of any finite sub-family is nonempty. So, suppose that $r \in \mathbb{N}$ and $\xi(x) = m_1 \otimes y_1 = \cdots = m_r \otimes y_r$ where $b_i \xrightarrow{m_i} a$ and $y_i \in Xb_i$. By [Lei1, 3.2] and an easy induction on r , there exist $c \xrightarrow{p} a$ and $z \in Xc$ such that $\xi(x) = p \otimes z$, and, for each $i \in \{1, \dots, r\}$, a map $g_i : c \longrightarrow b_i$ such that $m_i g_i = p$ and $g_i z = y_i$. But then for each i ,

$$\begin{aligned} \psi_p K_n(z) &= \psi_{m_i g_i} K_n(z) = \psi_{m_i}(Jg_i) K_n(z) \\ &\subseteq \psi_{m_i} K_n(g_i z) = \psi_{m_i} K_n(y_i), \end{aligned}$$

so

$$\psi_p K_n(z) \subseteq \psi_{m_1} K_n(y_1) \cap \cdots \cap \psi_{m_r} K_n(y_r),$$

and $\psi_p K_n(z) \neq \emptyset$ by inductive hypothesis.

Claim $|\bigcap_{n \in \mathbb{N}} K_n(x)| = 1$ for each $a \in \mathbb{A}$, $x \in Xa$.

Proof Let $a \in \mathbb{A}$ and $x \in Xa$. The subsets $K_n(x)$ of the compact space Ja are nonempty, closed, and form a nested sequence, so $\bigcap_{n \in \mathbb{N}} K_n(x)$ has at least one element. On the other hand, we may choose a resolution

$$(\cdots \xrightarrow{m_2} a_1 \xrightarrow{m_1} a_0, (x_n)_{n \in \mathbb{N}})$$

of x ; then for each $n \in \mathbb{N}$ we have $K_n(x) \subseteq \psi_{m_1} \cdots \psi_{m_n}(Ja_n)$, so by (c), $\bigcap_{n \in \mathbb{N}} K_n(x)$ has at most one element.

By the last claim, there is for each $a \in \mathbb{A}$ a function $\bar{\xi}_a : Xa \longrightarrow Ja$ defined by $\bigcap_{n \in \mathbb{N}} K_n(x) = \{\bar{\xi}_a(x)\}$.

Claim $(\bar{\xi}_a)_{a \in \mathbb{A}}$ is a natural transformation $X \longrightarrow J$ of **Set**-valued functors.

Proof Let $f : a \longrightarrow a'$ be a map in $\mathbb{A}$ and $x \in Xa$. Then for all $n \in \mathbb{N}$,

$$f\bar{\xi}_a(x) \in (Jf)K_n(x) \subseteq K_n(fx),$$

so $f\bar{\xi}_a(x) = \bar{\xi}_{a'}(fx)$, as required.

Claim $\bar{\xi} : (X, \xi) \longrightarrow (J, \gamma)$ is a map of coalgebras in **Set**.

Proof We have to show that for all $a \in \mathbb{A}$, the square

$$\begin{array}{ccc} Xa & \xrightarrow{\xi_a} & (M \otimes X)a \\ \bar{\xi}_a \downarrow & & \downarrow (M \otimes \bar{\xi})_a \\ Ja & \xrightarrow{\gamma_a} & (M \otimes J)a \end{array}$$

commutes. So let $x \in Xa$ and write $\xi_a(x) = (b \xrightarrow{m} a) \otimes y$; we have to show that $\gamma_a \bar{\xi}_a(x) = m \otimes \bar{\xi}_b(y)$, or equivalently, $\bar{\xi}_a(x) = \psi_m \bar{\xi}_b(y)$, or equivalently, $\psi_m \bar{\xi}_b(y) \in K_n(x)$ for all $n \in \mathbb{N}$. When $n = 0$ this is certainly true. Now let $n \in \mathbb{N}$; we have to show that if $\xi_a(x) = (b' \xrightarrow{m'} a) \otimes y'$ then $\psi_m \bar{\xi}_b(y) \in \psi_{m'} K_n(y')$. Since $m \otimes y = m' \otimes y'$, there exist a commutative square

$$\begin{array}{ccc} & c & \\ g \swarrow & & \searrow g' \\ b & & b' \\ m \searrow & & \swarrow m' \\ & a & \end{array}$$

and $z \in Xc$ such that $gz = y$ and $g'z = y'$. Hence

$$\begin{aligned}\psi_m \bar{\xi}_b(y) &= \psi_m \bar{\xi}_b(gz) = \psi_m g \bar{\xi}_c(z) \\ &= \psi_m g \bar{\xi}_c(z) = \psi_{m'g'} \bar{\xi}_c(z) \\ &= \psi_{m'} \bar{\xi}_{b'}(y') \in \psi_{m'} K_n(y')\end{aligned}$$

(the last equality by symmetry), as required.

Claim If $\tilde{\xi} : (X, \xi) \longrightarrow (J, \gamma)$ is a map of coalgebras in **Set** then $\tilde{\xi} = \bar{\xi}$.

Proof We show by induction on $n \in \mathbb{N}$ that $\tilde{\xi}_a(x) \in K_n(x)$ for all $a \in \mathbb{A}$ and $x \in Xa$; the result follows. For $n = 0$ this is trivial. Let $n \in \mathbb{N}$, $a \in \mathbb{A}$, and $x \in Xa$. If $\xi_a(x) = (b \xrightarrow{m} a) \otimes y$ then, as in the proof of the previous claim, $\tilde{\xi}$ being a map of coalgebras implies that $\tilde{\xi}_a(x) = \psi_m \tilde{\xi}_b(y)$, so by inductive hypothesis $\tilde{\xi}_a(x) \in \psi_m K_n(y)$. Hence $\tilde{\xi}_a(x) \in K_{n+1}(x)$, as required.

So (J, γ) (or properly, $U_*(J, \gamma)$) is the terminal coalgebra in **Set**. By Theorem A.1 of [Lei1], condition **S** holds, so $U_*(J, \gamma)$ is the universal solution $U_*(I, \iota)$ constructed in §4 of [Lei1]. There is a unique map $(J, \gamma) \longrightarrow (I, \iota)$ of coalgebras in **Top**, each component $Ja \longrightarrow Ia$ of which is a continuous bijection of compact Hausdorff spaces, hence a homeomorphism. So (J, γ) is isomorphic to (I, ι) as a coalgebra in **Top**, and is therefore the universal solution in **Top**. $\square$

In many examples the universal solution is especially easy to recognize.

Corollary 1.5 (Crude Recognition Theorem) *Suppose that $\mathbb{A}$ is finite. Let $(J, \gamma = \psi^{-1})$ be a fixed point of M in **Top** such that for each $a \in \mathbb{A}$, Ja is nonempty and compact, and suppose that the spaces Ja can be metrized in such a way that for each $b \xrightarrow{m} a$, the map $Jb \xrightarrow{\psi_m} Ja$ is a contraction. Then (J, γ) is the universal solution of $(\mathbb{A}, M)$.*

Proof Since $\mathbb{A}$ and M are finite, there are only finitely many module elements m , so we may choose $\lambda < 1$ such that each ψ_m is a contraction with constant λ . Since $\mathbb{A}$ is finite and each Ja compact, we may also choose $D \geq 0$ such that $\text{diam}(Ja) \leq D$ for all $a \in \mathbb{A}$.

We verify condition (b) of the Precise Recognition Theorem. Certainly J is occupied. For the main part of the condition, let $\varepsilon > 0$; then we may choose $n \in \mathbb{N}$ such that $\lambda^n D \leq \varepsilon$, and the result follows. $\square$

2 Examples

Example 2.1 (Interval) We finally prove the Topological Freyd Theorem [Lei1, 1.7]. So far we have verified that the $(\mathbb{A}, M)$ concerned is a self-similarity system and exhibited an M -coalgebra (J, γ) (previously written

(I, ι) with $J(1) = [0, 1]$. Evidently γ is invertible, so we have a fixed point $(J, \gamma = \psi^{-1})$. The only non-constant maps ψ_m are

$$\begin{array}{ccc} [0, 1] & \longrightarrow & [0, 1] \\ t & \longmapsto & t/2 \\ t & \longmapsto & (t+1)/2, \end{array}$$

both of which are contractions, so by the Crude Recognition Theorem, (J, γ) is the universal solution.

Freyd's Theorem expresses $[0, 1]$ as two copies of itself glued end to end. Two can be replaced by any larger number. Thus, for each $k \geq 2$ there is a self-similarity structure $M^{(k)}$ on $\mathbb{A}$ (with, for instance, $|M^{(k)}(1, 1)| = k$), the multiplication map $k \cdot - : [0, 1] \longrightarrow [0, k]$ puts an $M^{(k)}$ -coalgebra structure $\gamma^{(k)}$ on J , and the same argument shows that $(J, \gamma^{(k)})$ is the universal solution.

Example 2.2 (Circle) In most of our examples, every arrow in $\mathbb{A}$ is monic. Here is an exception. Let $\mathbb{A}$ be the category generated by objects and arrows

$$0 \xrightleftharpoons[\tau]{\sigma} 1 \xrightarrow{\rho} 2$$

subject to $\rho\sigma = \rho\tau$, as in [Lei1, 2.12]. There is a self-similarity structure M on $\mathbb{A}$ given informally by the equations

$$\begin{array}{c} X_0 = X_0 \\ \begin{array}{c} \text{Diagram 1: } X_1 \text{ containing two } X_0 \text{ objects.} \\ \text{Diagram 2: } X_1 \text{ containing three } X_0 \text{ objects.} \end{array} \\ X_2 = \begin{array}{c} \text{Diagram 3: } X_1 \text{ containing } X_0 \text{ at the top and bottom, with } X_1 \text{ in the middle.} \end{array} \end{array}$$

and formally by

$$M(-, 0) \xrightleftharpoons[\tau \cdot -]{\sigma \cdot -} M(-, 1) \xrightarrow{\rho \cdot -} M(-, 2)$$

$$\begin{array}{ccccc} M(0, -) & \{ \text{id} \} & \xrightleftharpoons[1]{0} & \{0, \frac{1}{2}, 1\} & \xrightarrow{\text{reduce mod } 1} & \{0, \frac{1}{2}\} \\ \uparrow \neg \sigma & \uparrow & & \uparrow \text{inf} & & \uparrow \text{inf} \\ \uparrow & & & & & \uparrow \\ M(1, -) & \emptyset & \xrightleftharpoons{\quad} & \{[0, \frac{1}{2}], [\frac{1}{2}, 1]\} & \xrightarrow{\text{id}} & \{[0, \frac{1}{2}], [\frac{1}{2}, 1]\} \\ \uparrow \neg \rho & \uparrow & & \uparrow & & \uparrow \\ M(2, -) & \emptyset & \xrightleftharpoons{\quad} & \emptyset & \longrightarrow & \emptyset. \end{array}$$

(The top-left quarter is the Freyd module: eq. (9) of [Lei1].) By [Lei1, 2.12], M is nondegenerate, and clearly M is finite, so $(\mathbb{A}, M)$ is a self-similarity system. We also have a nondegenerate functor $J : \mathbb{A} \longrightarrow \mathbf{Top}$ given by

$$\{\star\} \xrightleftharpoons[1]{0} [0, 1] \xrightarrow{q} S^1$$

where q identifies the endpoints. Now $M \otimes J$ can naturally be identified with

$$\{\star\} \xrightleftharpoons[2]{0} [0, 2] \xrightarrow{\text{quotient}} [0, 2]/(0 = 2),$$

so there is an obvious isomorphism $\gamma : J \xrightarrow{\sim} M \otimes J$. Moreover, each of the spaces Ja is compact and nonempty. The non-trivial maps of the form ψ_m (where $\psi = \gamma^{-1}$) are the two maps

$$\psi_{[0, \frac{1}{2}]}, \psi_{[\frac{1}{2}, 1]} : [0, 1] \longrightarrow [0, 1]$$

seen in Example 2.1 and their respective composites with $q : [0, 1] \longrightarrow S^1$. If we give $[0, 1]$ the standard metric and S^1 the standard metric scaled down by a sufficiently large factor then all four maps are contractions, so by the Crude Recognition Theorem, (J, γ) is the universal solution.

Example 2.3 (Cantor set) Write $\mathbf{1}$ for the terminal category (one object and only the identity arrow). A self-similarity structure on $\mathbf{1}$ is a finite set M , and an M -coalgebra is a space X equipped with a map into the M -fold coproduct $M \times X$. The universal solution is the power $M^{\mathbb{N}}$ (regarding M as a discrete space) together with the shift isomorphism $\gamma = \psi^{-1} : M^{\mathbb{N}} \xrightarrow{\sim} M \times M^{\mathbb{N}}$. This can be seen directly from the construction of the universal solution in [Lei1, §4], or from a Recognition Theorem as follows. The space $M^{\mathbb{N}}$ is nonempty if M is, hence occupied. It is compact. For $m \in M$, the map $\psi_m : M^{\mathbb{N}} \longrightarrow M^{\mathbb{N}}$ is given by

$$\psi_m(m_0, m_1, \dots) = (m, m_0, m_1, \dots),$$

so condition (c) of the Precise Recognition Theorem holds, as required. When M has cardinality 2, the universal solution is the standard Cantor set $\{0, 1\}^{\mathbb{N}}$, often regarded as a subspace of $[0, 1]$ via the embedding $(m_n)_{n \in \mathbb{N}} \longmapsto \sum_{n \in \mathbb{N}} 2m_n \cdot 3^{-(n+1)}$.

In fact, the homeomorphism type of $M^{\mathbb{N}}$ is independent of M for $|M| \geq 2$. This can be proved directly or as follows. Write $k = \{0, \dots, k-1\}$, write $\psi : 2 \times 2^{\mathbb{N}} \xrightarrow{\sim} 2^{\mathbb{N}}$ for the (inverse) shift isomorphism, and define for each $k \geq 1$ an isomorphism $\psi^{(k)} : k \times 2^{\mathbb{N}} \xrightarrow{\sim} 2^{\mathbb{N}}$ by

- $\psi^{(1)}$ is the canonical isomorphism
- $\psi^{(k+1)} = \left((k+1) \times 2^{\mathbb{N}} \xrightarrow[\sim]{\text{canonical}} (k \times 2^{\mathbb{N}}) + 2^{\mathbb{N}} \xrightarrow[\sim]{\psi^{(k)} + 1} 2^{\mathbb{N}} + 2^{\mathbb{N}} \xrightarrow[\sim]{\psi} 2^{\mathbb{N}} \right).$

Then for each $k \geq 1$ and $m \in k$, the map $\psi_m^{(k)} : 2^{\mathbb{N}} \longrightarrow 2^{\mathbb{N}}$ is of the form $\psi_{p_1} \cdots \psi_{p_r}$ with $r \geq 0$ and $p_j \in 2$; if $k \geq 2$ then $r \geq 1$. Using the metric on $2^{\mathbb{N}}$ induced by its embedding into $[0, 1]$, ψ_0 and ψ_1 are contractions with constant $1/3$, so if $k \geq 2$ then each $\psi_m^{(k)}$ is also a contraction with constant $1/3$. So by the Crude Recognition Theorem, $(2^{\mathbb{N}}, (\psi^{(k)})^{-1})$ is the universal solution of $(\mathbf{1}, k)$ whenever $k \geq 2$. In particular, $2^{\mathbb{N}} \cong k^{\mathbb{N}}$.

Example 2.4 (Discrete systems) In §4 we will look systematically at self-similarity systems $(\mathbb{A}, M)$ in which $\mathbb{A}$ is a discrete category. Proposition 4.9 says that the universal solution often consists simply of copies of the Cantor set and $\emptyset$. But it is not always so: for instance, the universal solution of the system

$$\begin{aligned} A &= A \\ B &= A + B \end{aligned}$$

(defined formally before eq. (6) of [Lei1]) is $A = \{0\}$ and $B = \mathbb{N} \cup \{\infty\}$, with B topologized as the Alexandroff compactification of the discrete space $\mathbb{N}$.

Finite products of self-similarity systems can be formed in the expected way.

Lemma 2.5 *Let $(\mathbb{B}_\omega \xrightarrow{Z_\omega} \mathbf{Set})_{\omega \in \Omega}$ be a family of functors on small categories $\mathbb{B}_\omega$, indexed over an arbitrary set Ω . Then there is a functor $\prod_{\omega \in \Omega} Z_\omega : \prod_{\omega \in \Omega} \mathbb{B}_\omega \longrightarrow \mathbf{Set}$, and*

- a. $\mathbb{E}(\prod Z_\omega) \cong \prod \mathbb{E}(Z_\omega)$
- b. if Ω is finite and each $\mathbb{E}(Z_\omega)$ is finite then so is $\mathbb{E}(\prod Z_\omega)$
- c. if each Z_ω is nondegenerate then so is $\prod Z_\omega$.

Proof (a) is straightforward, and (b) and (c) follow. $\square$

Lemma 2.6 *Let $(\mathbb{B}_\omega \xrightarrow{X_\omega} \mathbf{Top})_{\omega \in \Omega}$ and $(\mathbb{B}_\omega^{\text{op}} \xrightarrow{Y_\omega} \mathbf{Set})_{\omega \in \Omega}$ be families of functors on small categories $\mathbb{B}_\omega$, where Ω is finite and each $X_\omega b$ is compact Hausdorff. Then*

$$\left(\prod_{\omega \in \Omega} Y_\omega \right) \otimes \left(\prod_{\omega \in \Omega} X_\omega \right) \cong \prod_{\omega \in \Omega} (Y_\omega \otimes X_\omega).$$

Proof This is clear when $\Omega = \emptyset$, so by induction, it suffices to consider $\Omega = \{1, 2\}$. In that case,

$$\begin{aligned} (Y_1 \times Y_2) \otimes (X_1 \times X_2) &\cong \int^{(b_1, b_2)} Y_1 b_1 \times Y_2 b_2 \times X_1 b_1 \times X_2 b_2 \\ &\cong \int^{b_1} Y_1 b_1 \times X_1 b_1 \times \int^{b_2} Y_2 b_2 \times X_2 b_2 \\ &\cong (Y_1 \otimes X_1) \times (Y_2 \otimes X_2), \end{aligned}$$

using in the second isomorphism the fact that if K is compact Hausdorff then $K \times - : \mathbf{Top} \longrightarrow \mathbf{Top}$ preserves colimits. $\square$

Proposition 2.7 (Product self-similarity system) *Let $((\mathbb{A}_\lambda, M_\lambda))_{\lambda \in \Lambda}$ be a finite family of self-similarity systems. Then there is a product self-similarity system $(\prod_{\lambda \in \Lambda} \mathbb{A}_\lambda, \prod_{\lambda \in \Lambda} M_\lambda)$, where*

$$\left(\prod_{\lambda \in \Lambda} M_\lambda \right) ((b_\lambda)_{\lambda \in \Lambda}, (a_\lambda)_{\lambda \in \Lambda}) = \prod_{\lambda \in \Lambda} M_\lambda(b_\lambda, a_\lambda).$$

If $(\mathbb{A}_\lambda, M_\lambda)$ has a universal solution $(I_\lambda, \iota_\lambda)$ for each $\lambda \in \Lambda$ then $(\prod \mathbb{A}_\lambda, \prod M_\lambda)$ has a universal solution $(\prod I_\lambda, \prod \iota_\lambda)$.

Proof Lemma 2.5 implies that $(\prod \mathbb{A}_\lambda, \prod M_\lambda)$ is a self-similarity system. Now suppose that each $(\mathbb{A}_\lambda, M_\lambda)$ has a universal solution $(I_\lambda, \iota_\lambda)$ in **Top**. The functor $\prod I_\lambda : \prod \mathbb{A}_\lambda \longrightarrow \mathbf{Set}$ is nondegenerate, since

$$U \circ (\prod I_\lambda) \cong \prod (U \circ I_\lambda) : \prod \mathbb{A}_\lambda \longrightarrow \mathbf{Set}$$

is nondegenerate by Lemma 2.5(c) and each space $(\prod_{\lambda \in \Lambda} I_\lambda)(a_\lambda)_{\lambda \in \Lambda}$ is compact Hausdorff. By Lemma 2.6, we have a coalgebra structure

$$\prod \iota_\lambda : \prod I_\lambda \longrightarrow \prod (M_\lambda \otimes I_\lambda) \cong (\prod M_\lambda) \otimes (\prod I_\lambda)$$

on $\prod I_\lambda$, and $\prod \iota_\lambda$ is an isomorphism. Also, $\prod I_\lambda$ is occupied since each I_λ is. To finish the proof it remains only to verify that $(\prod I_\lambda, \prod \iota_\lambda)$ satisfies the main condition in (c) of the Precise Recognition Theorem, and this follows from the fact that it is satisfied by each $(I_\lambda, \iota_\lambda)$. $\square$

Example 2.8 (Cubes) Let $(\mathbb{A}, M)$ be the Freyd self-similarity system. Then by Proposition 2.7, the universal solution of $(\mathbb{A}^2, M^2)$ has a universal solution (I, ι) satisfying $I(1, 1) = [0, 1]^2$. Informally, the self-similarity equations are

The diagrammatic equations are as follows:

- A single point $\bullet$ is equal to a single point $\bullet$.
- A horizontal line segment with endpoints $\bullet$ is equal to a horizontal line segment with endpoints $\bullet$.
- A vertical line segment with endpoints $\bullet$ is equal to a vertical line segment with endpoints $\bullet$.
- A square with vertices $\bullet$ is equal to a larger square divided into four smaller squares, each with vertices $\bullet$.



For general $n \in \mathbb{N}$, $(\mathbb{A}^n, M^n)$ has universal solution given by $[0, 1]^n$.

Example 2.9 Similarly, if C denotes the Cantor set then the space

$$C \times [0, 1] = \begin{array}{c} \text{||||} \\ \text{||||} \end{array}$$

[illegible]

Example 2.10 (Sierpiński simplices) Let $n \in \mathbb{N}$ and let $s_0, \dots, s_n$ be affinely independent points of $\mathbb{R}^n$. For each $i \in \{0, \dots, n\}$, write $\psi_i : \mathbb{R}^n \rightarrow \mathbb{R}^n$ for the scaling with fixed point s_i and scale factor $\frac{1}{2}$. The **Sierpiński simplex** with vertices $s_0, \dots, s_n$ is the unique nonempty compact subset S of $\mathbb{R}^n$ satisfying $S = \bigcup_{i=0}^n \psi_i S$. When $n = 1$, this is the closed interval with endpoints s_0 and s_1 . When $n = 2$, this is the usual Sierpiński triangle or gasket, which satisfies the equation


$$M(-, 0) \begin{array}{c} \xrightarrow{\sigma_0 \cdot -} \\ \vdots \\ \xrightarrow{\sigma_n \cdot -} \end{array} M(-, 1)$$


Then $(\mathbb{A}, M)$ is a self-similarity system. Any space X equipped with distinct basepoints $x_0, \dots, x_n$ defines a nondegenerate functor $X : \mathbb{A} \longrightarrow \mathbf{Top}$, and then $M \otimes X$ is the quotient space

$$\frac{\{0, \dots, n\} \times X}{(i, x_j) = (j, x_i) \text{ for all } i, j}$$

with basepoints $(0, x_0), \dots, (n, x_n)$. Choose affinely independent points $s_0, \dots, s_n \in \mathbb{R}^n$ and take S as above. Then $(S, s_0, \dots, s_n)$ determines a nondegenerate functor $J : \mathbb{A} \longrightarrow \mathbf{Top}$, and $\psi_0, \dots, \psi_n$ determine a basepoint-preserving map

$$\begin{array}{ccc} \{0, \dots, n\} \times S & \longrightarrow & S \\ (i, s) & \longmapsto & \psi_i(s). \end{array} \quad (1)$$

If $i \neq j$ then the only point in $\psi_i S \cap \psi_j S$ is the midpoint $\psi_i s_j = \psi_j s_i$ of s_i and s_j . Also, each ψ_i is injective and, by definition of S , the map (1) is surjective, so (1) induces a basepoint-preserving homeomorphism $(M \otimes J)1 \xrightarrow{\sim} J1$. In other words, we have an isomorphism $\psi : M \otimes J \xrightarrow{\sim} J$. The spaces $J0 = \{\star\}$ and $J1 = S$ are nonempty and compact and the structure maps $\psi_i : J1 \longrightarrow J1$ are contractions, so by the Crude Recognition Theorem, (J, ψ^{-1}) is the universal solution of $(\mathbb{A}, M)$.

Example 2.11 (IFSs) More generally, let $(\psi_0, \dots, \psi_n)$ be an iterated function system on $\mathbb{R}^d$ (some $d \in \mathbb{N}$); write S for the unique nonempty compact subset of $\mathbb{R}^d$ satisfying $S = \bigcup_{i=0}^n \psi_i S$, and s_i for the fixed point of ψ_i . Suppose that $\psi_0, \dots, \psi_n$ are injective, that $s_0, \dots, s_n$ are distinct, and that if $\psi_i s = \psi_j t$ with $i \neq j$ and $s, t \in S$ then $s, t \in \{s_0, \dots, s_n\}$. Then the space S arises from a self-similarity system, as follows.

Define an equivalence relation $\sim$ on $\{0, \dots, n\}^2$ by $(i, j) \sim (i', j') \iff \psi_i s_j = \psi_{i'} s_{j'}$, and write $[i, j]$ for the equivalence class of (i, j) . Exactly as in the previous example, this equivalence relation gives rise to a self-similarity system $(\mathbb{A}, M)$, the space S with basepoints $s_0, \dots, s_n$ determines a nondegenerate functor $J : \mathbb{A} \longrightarrow \mathbf{Top}$, the maps ψ_i determine an isomorphism $M \otimes J \xrightarrow{\sim} J$, and by the Crude Recognition Theorem, this is the universal solution of $(\mathbb{A}, M)$.

Solutions to iterated function systems in which the small copies $\psi_i S$ of S overlap by more than just a point can sometimes be described as universal solutions too: $[0, 1]^n$ in Example 2.8, for instance.

Example 2.12 (Barycentric subdivision) Barycentric subdivision expresses the n -simplex Δ^n as $(n+1)!$ smaller copies of itself glued together along simplices of lower dimension. This self-similarity can be formalized as follows.

Let Δ_{face} be the category whose objects are the nonempty finite totally ordered sets $[n] = \{0, \dots, n\}$ ($n \in \mathbb{N}$) and whose maps are the order-preserving injections. For each $n, m \in \mathbb{N}$, put

$$M([n], [m]) = \{\text{chains } \emptyset \subset Q(0) \subset \dots \subset Q(n) \subseteq [m]\}$$




Figure 2-A: Barycentric subdivision of Δ^2

where $\subset$ means proper subset. The idea can be seen in Figure 2-A: the 1-simplex in bold and the shaded 2-simplex correspond respectively to

$$\begin{aligned} (\emptyset \subset \{0, 2\} \subset \{0, 1, 2\}) &\in M([1], [2]), \\ (\emptyset \subset \{0\} \subset \{0, 2\} \subset \{0, 1, 2\}) &\in M([2], [2]). \end{aligned}$$

An element of $M([n], [m])$ can be regarded as an order-preserving injection $[n] \longrightarrow \mathcal{P}_{\neq \emptyset}[m]$, where $\mathcal{P}_{\neq \emptyset}$ denotes the set of nonempty subsets ordered by inclusion. By using direct images, $\mathcal{P}_{\neq \emptyset}[m]$ is functorial in $[m]$, so M defines a module $\Delta_{\text{face}} \dashrightarrow \Delta_{\text{face}}$. It is straightforward to show that Δ_{face} has all pullbacks and equalizers and that $\mathcal{P}_{\neq \emptyset} : \Delta_{\text{face}} \longrightarrow \mathbf{Poset}$ preserves them, from which it follows that M is nondegenerate [Lei1, 2.9]. And clearly M is finite, so $(\Delta_{\text{face}}, M)$ is a self-similarity system.

We are going to show that the universal solution is given by the simplex functor $\Delta^\bullet : \Delta_{\text{face}} \longrightarrow \mathbf{Top}$. To define $\Delta^\bullet$, choose for each $n \in \mathbb{N}$ an affinely independent sequence $e_0^n, \dots, e_n^n$ of elements of $\mathbb{R}^n$ and let Δ^n be their convex hull; then for each map $f : [n] \longrightarrow [m]$ in Δ_{face} there is a unique affine map $\mathbb{R}^n \longrightarrow \mathbb{R}^m$ sending e_j^n to $e_{f(j)}^m$ for each j , which restricts to a map $\Delta f = f_* : \Delta^n \longrightarrow \Delta^m$. Again it is straightforward to check that $U \circ \Delta^\bullet : \Delta_{\text{face}} \longrightarrow \mathbf{Set}$ preserves pullbacks and equalizers, and each space Δ^n is compact Hausdorff, so $\Delta^\bullet$ is nondegenerate.

Given $Q : [n] \dashrightarrow [m]$, there is a unique affine map $\mathbb{R}^n \longrightarrow \mathbb{R}^m$ such that

$$e_j^n \mapsto \frac{1}{|Q(j)|} \sum_{i \in Q(j)} e_i^m$$

for all $j \in [n]$, and this restricts to a map $\psi_Q : \Delta^n \longrightarrow \Delta^m$. It can be checked that $\psi_{fQg} = f_* \circ \psi_Q \circ g_*$ for all f, Q , and g , so by the remarks at the beginning of §1, ψ defines an M -algebra structure on $\Delta^\bullet$.

A standard calculation [Hat, 2.21] shows that for any $Q : [n] \dashrightarrow [m]$,

$$\text{diam}(\psi_Q \Delta^n) \leq \frac{m}{m+1} \text{diam}(\Delta^m)$$

in the Euclidean metric. More generally, if

$$[n_r] \xrightarrow{Q_r} \cdots \xrightarrow{Q_1} [n_0]$$

then the same method shows that

$$\begin{aligned} \text{diam}(\psi_{Q_1} \cdots \psi_{Q_r} \Delta^{n_r}) &\leq \left(\frac{n_{r-1}}{n_{r-1} + 1} \right) \cdots \left(\frac{n_0}{n_0 + 1} \right) \text{diam}(\Delta^{n_0}) \\ &\leq \left(\frac{n_0}{n_0 + 1} \right)^r \text{diam}(\Delta^{n_0}). \end{aligned}$$

So as long as ψ is an isomorphism, condition (b) of the Precise Recognition Theorem holds and $(\Delta^\bullet, \psi^{-1})$ is the universal solution.

It remains to show that ψ is an isomorphism. Since the spaces involved are compact Hausdorff, this says that for each $m \in \mathbb{N}$,

$$\psi : \int^{[n]} M([n], [m]) \times \Delta^n \longrightarrow \Delta^m$$

is a bijection. The proof is somewhat technical.

For surjectivity, take $s \in \Delta^m$; then $s = \sum_{i=0}^m \sigma_i e_i^m$ with $\sigma_i \geq 0$ and $\sum \sigma_i = 1$. There are unique $n \in \mathbb{N}$ and $\kappa_0 > \cdots > \kappa_n > \kappa_{n+1} = 0$ such that

$$\{\kappa_0, \dots, \kappa_n, \kappa_{n+1}\} = \{\sigma_0, \dots, \sigma_m, 0\},$$

and we may define $q : \{0, \dots, m\} \longrightarrow \{0, \dots, n+1\}$ by $\kappa_{q(i)} = \sigma_i$. Put

$$Q(j) = q^{-1}\{0, \dots, j\}, \quad \tau_j = (\kappa_j - \kappa_{j+1})|Q(j)|, \quad t = \sum_{j=0}^n \tau_j e_j^n.$$

Then $Q \in M([n], [m])$, $t \in \Delta^n$, and $\psi_Q(t) = s$, as required.

For injectivity, let $s \in \Delta^m$ and take $Q' : [n'] \dashrightarrow [m]$ and $t' \in \Delta^{n'}$ with $\psi_{Q'}(t') = s$. It will be enough to show that with $[n]$, Q and t as in the previous paragraph, there exists $f : [n] \longrightarrow [n']$ such that

$$\begin{array}{ccc} [n] & & \\ & \searrow Q & \\ & & [m] \\ & \nearrow Q' & \\ [n'] & & \end{array}$$

commutes and $f_* t = t'$. A map $f : [n] \longrightarrow [n']$ in Δ_{face} amounts to an $(n+1)$ -element subset of $[n']$, and it can be seen that $\{j' \in [n'] \mid t'_{j'} \neq 0\}$ has $n+1$ elements. This defines f ; the remaining checks are left to the reader.

Example 2.13 (Edgewise subdivision) Edgewise subdivision (Figure 2-B and [Freu]) expresses Δ^n as 2^n smaller copies of itself glued together. This too can be formalized by a self-similarity structure M on Δ_{face} . Here,

$$\begin{aligned} M([n], [m]) &= \{\text{order-preserving injections } (p, q) : [n] \longrightarrow [m] \times [m] \\ &\quad \text{such that } p(n) \leq q(0)\} \end{aligned}$$

where $[m] \times [m]$ is the product in the category of posets. For instance, the shaded 2-simplex in Figure 2-B is the module element $[2] \dashrightarrow [3]$ given by the order-preserving injection $[2] \longrightarrow [3] \times [3]$ with image $\{(0, 2), (0, 3), (1, 3)\}$. Again it can be shown that $\Delta^\bullet : \Delta_{\text{face}} \longrightarrow \mathbf{Top}$, equipped with a certain M -coalgebra structure, is the universal solution.




Figure 2-B: Edgewise subdivisions of Δ^2 and Δ^3

3 Spaces admitting a self-similarity

Here we answer the question: which topological spaces admit at least one self-similarity structure? That is, call a space **self-similar** if it is homeomorphic to Ia for some self-similarity system $(\mathbb{A}, M)$ with universal solution (I, ι) and some $a \in \mathbb{A}$; which spaces are self-similar? We know that every self-similar space is compact metrizable. In fact, the converse holds:

Theorem 3.1 (Self-similar spaces) *A topological space is self-similar if and only if it is compact and metrizable.*

This calls for some explanation.

First recall that when we formalized the notion of self-similarity system [Lei1, §1], we allowed there to be infinitely many equations, even though each individual equation could involve only finitely many spaces. So there can be infinite regress: X_1 might be described as a copy of itself copied to a copy of X_2 , X_2 as a copy of itself glued to a copy of X_3 , and so on. This is essential to the proof of Theorem 3.1. The more restrictive notion of finite self-similarity is discussed at the end of the section.

Second, this theorem does not exhaust the subject. It characterizes those spaces that are self-similar in at least one way, but the same space may carry several different self-similarities: for instance, the spaces Δ^n are self-similar by both barycentric and edgewise subdivision (§2). Compare the result that every nonempty set admits at least one group structure, which does not exhaust group theory.

Here is the strategy for the proof. Let S be a compact metrizable space. Cover S by two closed subsets V_1 and V'_1 . Then $S = V_1 \cup V'_1$; more precisely, S is the pushout

$$S = V_1 +_{V_1''} V'_1$$

where $V_1'' = V_1 \cap V'_1$. Next, cover S by a different pair V_2, V'_2 of closed subsets and write $V_2'' = V_2 \cap V'_2$: then

$$\begin{aligned} V_1 &= (V_1 \cap V_2) +_{(V_1 \cap V_2'')} (V_1 \cap V_2'), \\ V'_1 &= (V'_1 \cap V_2) +_{(V'_1 \cap V_2'')} (V'_1 \cap V_2'), \\ V_1'' &= (V_1'' \cap V_2) +_{(V_1'' \cap V_2'')} (V_1'' \cap V_2'). \end{aligned}$$

Continue in this way to obtain a countable self-similarity system. Compact metrizable of S means that the covers can be chosen to penetrate all of its structure, and the universal solution I is then made up of the spaces $S, V_1, V'_1, V''_1, V_2, V'_2, V''_2, \dots$ and inclusions between them.

Formally, a **separating sequence** on a space S is a sequence $(\mathcal{V}_n)_{n \geq 1}$ of finite closed covers of S such that

- if $n \geq 1$ and $V, V' \in \mathcal{V}_n$ then $V \cap V' \in \mathcal{V}_n$
- if $s, s' \in S$ and for all $n \geq 1$ there exists $V \in \mathcal{V}_n$ containing both s and s' then $s = s'$.

Lemma 3.2 *Every compact metrizable space admits a separating sequence.*

Proof Every compact metrizable space S has a basis $(U_n)_{n \geq 1}$ of open sets. For each $n \geq 1$, let

$$V_n = \text{closure}(U_n), \quad V'_n = S \setminus U_n, \quad V''_n = V_n \cap V'_n,$$

and $\mathcal{V}_n = \{V_n, V'_n, V''_n\}$. Then $(\mathcal{V}_n)_{n \geq 1}$ is a sequence of finite closed covers satisfying the binary intersection axiom. Let s and s' be distinct points of S . Since S is Hausdorff, there exists $n \geq 1$ with $s \in U_n$ but $s' \notin V_n$; then there is no $V \in \mathcal{V}_n$ such that $s, s' \in V$. $\square$

Fix a compact metrizable space S with a separating sequence $(\mathcal{V}_n)_{n \geq 1}$.

First we define a self-similarity system $(\mathbb{A}, M)$. Recall that a poset can be regarded as a category in which each hom-set has at most one element: there is a map $a' \longrightarrow a$ just when $a' \leq a$. Let $\mathbb{A} = \sum_{n \geq 0} \mathbb{A}_n$, where $\mathbb{A}_n$ is the sub-poset of $(\mathcal{V}_1, \subseteq) \times \dots \times (\mathcal{V}_n, \subseteq)$ consisting of those $(V_1, \dots, V_n)$ for which $V_1 \cap \dots \cap V_n \neq \emptyset$. The module M is also ‘posetal’, that is, $M(b, a)$ never has more than one element; there is a module element

$$(W_1, \dots, W_m) \dashrightarrow (V_1, \dots, V_n)$$

if and only if $m = n + 1$ and $W_1 \cap \dots \cap W_{n+1} \subseteq V_1 \cap \dots \cap V_n$.

Lemma 3.3 $(\mathbb{A}, M)$ is a self-similarity system.

Proof M is finite since each $\mathcal{V}_n$ is finite. For nondegeneracy, the binary intersection axiom guarantees that the poset $\mathbb{A}$ has all binary meets. Hence the category $\mathbb{A}$ has all pullbacks; they are the squares in $\mathbb{A}$ of the form

$$\begin{array}{ccc} & (V_1 \cap V'_1, \dots, V_n \cap V'_n) & \\ \swarrow & & \searrow \\ (V_1, \dots, V_n) & & (V'_1, \dots, V'_n) \\ \searrow & & \swarrow \\ & (W_1, \dots, W_n) & \end{array}$$

So by [Lei1, 2.9], a functor $X : \mathbb{A} \longrightarrow \mathbf{Set}$ is nondegenerate just when it preserves pullbacks (existence and preservation of equalizers being trivial). This is easily verified when $X = M(b, -)$ for some $b \in \mathbb{A}$. $\square$

Next we define an M -algebra (J, ψ) in **Top**. The functor $J : \mathbb{A} \longrightarrow \mathbf{Top}$ is given by

$$J(V_1, \dots, V_n) = V_1 \cap \dots \cap V_n \subseteq S,$$

and if $V_1 \subseteq V'_1, \dots, V_n \subseteq V'_n$ then J sends the resulting arrow in $\mathbb{A}$ to the inclusion $V_1 \cap \dots \cap V_n \hookrightarrow V'_1 \cap \dots \cap V'_n$. The map $\psi : M \otimes J \longrightarrow J$ is defined by taking its component at a module element

$$(W_1, \dots, W_{n+1}) \dashrightarrow (V_1, \dots, V_n),$$

to be the inclusion

$$W_1 \cap \dots \cap W_{n+1} \hookrightarrow V_1 \cap \dots \cap V_n.$$

Lemma 3.4 $J : \mathbb{A} \longrightarrow \mathbf{Top}$ is a nondegenerate functor and $\psi : M \otimes J \longrightarrow J$ is an isomorphism.

Proof For the first part, each space $J(V_1, \dots, V_n)$ is compact Hausdorff, so it suffices to show that $U \circ J : \mathbb{A} \longrightarrow \mathbf{Set}$ is nondegenerate; and as in the proof of 3.3, this reduces to a straightforward check that $U \circ J$ preserves pullbacks.

We now have to show that for each $(V_1, \dots, V_n) \in \mathbb{A}$, the canonical map

$$\psi : \int_{\substack{(W_1, \dots, W_{n+1}) \in \mathbb{A}_{n+1} : \\ W_1 \cap \dots \cap W_{n+1} \subseteq V_1 \cap \dots \cap V_n}} W_1 \cap \dots \cap W_{n+1} \longrightarrow V_1 \cap \dots \cap V_n$$

is a homeomorphism. The domain is a finite colimit of compact spaces, hence compact, and the codomain is Hausdorff, so it suffices to show that ψ is a bijection.

For surjectivity, let $s \in V_1 \cap \dots \cap V_n$. We have $s \in V_{n+1}$ for some $V_{n+1} \in \mathcal{V}_{n+1}$, so then $s \in V_1 \cap \dots \cap V_{n+1}$ with $(V_1, \dots, V_{n+1}) \in \mathbb{A}_{n+1}$, and ψ sends $s \in V_1 \cap \dots \cap V_{n+1}$ to $s \in V_1 \cap \dots \cap V_n$.

For injectivity, suppose that

$$t \in W_1 \cap \dots \cap W_{n+1} \subseteq V_1 \cap \dots \cap V_n, \quad t' \in W'_1 \cap \dots \cap W'_{n+1} \subseteq V_1 \cap \dots \cap V_n$$

with $\psi(t) = \psi(t')$; then t and t' are equal as elements of S . By the binary intersection property there is a diagram

$$\begin{array}{ccc} & (W_1 \cap W'_1, \dots, W_{n+1} \cap W'_{n+1}) & \\ \swarrow & & \searrow \\ (W_1, \dots, W_{n+1}) & & (W'_1, \dots, W'_{n+1}) \end{array} \quad (2)$$

in $\mathbb{A}$ and an element

$$t'' \in (W_1 \cap W'_1) \cap \dots \cap (W_{n+1} \cap W'_{n+1})$$

equal to t and t' as an element of S . By (2), t'' represents the same element of the coend as both t and t' , so t and t' represent the same element of the coend, as required. $\square$

Proposition 3.5 (J, ψ^{-1}) is the universal solution of $(\mathbb{A}, M)$ in **Top**.

Proof Each space $J(V_1, \dots, V_n)$ is compact, and nonempty by definition of $\mathbb{A}$, so it only remains to verify the main part of condition (c) of the Precise Recognition Theorem. Let $a = (V_1, \dots, V_n) \in \mathbb{A}$, let $t, t' \in V_1 \cap \dots \cap V_n$, and suppose there exists $(\dots \xrightarrow{m_1} a_0) \in \mathcal{I}a$ such that $t, t' \in \psi_{m_1} \dots \psi_{m_k}(Ja_k)$ for all $k \in \mathbb{N}$. Writing $a_k = (V_1^k, \dots, V_{n+k}^k) \in \mathbb{A}_{n+k}$, we have $t, t' \in V_1^k \cap \dots \cap V_{n+k}^k \subseteq V_k^k \in \mathcal{V}_k$ for all $k \geq 1$, so $t = t'$ by definition of separating sequence. $\square$

Theorem 3.1 follows: for if S is nonempty, the sequence $()$ of length 0 is an object of $\mathbb{A}$, and $J() = S$; on the other hand, the empty space is the universal solution of the self-similarity system $(\mathbf{1}, \emptyset)$ (Example 2.3).

The theorem has a restricted form, as follows.

A self-similarity system $(\mathbb{A}, M)$ is **discrete** if the category $\mathbb{A}$ is discrete (has no arrows except identities). Any **Set**-valued functor on a discrete category is nondegenerate, so a discrete self-similarity system consists of a set $\mathbb{A}$ and a family $(M(b, a))_{b, a \in \mathbb{A}}$ of sets such that $\sum_b M(b, a)$ is finite for each $a \in \mathbb{A}$. Condition **S** always holds, so every discrete self-similarity system has a universal solution: it is $(\text{ob } \mathcal{I}, \iota)$, where $\text{ob } \mathcal{I} : \mathbb{A} \longrightarrow \mathbf{Top}$ is the functor constructed before Lemma 6.5 of [Leil]. To see this, just note that for each $a \in \mathbb{A}$, the canonical map $\pi_a : \text{ob } (\mathcal{I}a) \longrightarrow Ia$ is a bijection of compact Hausdorff spaces, hence a homeomorphism.

A **discretely self-similar** space is one homeomorphic to Ia for some discrete self-similarity system $(\mathbb{A}, M)$ with universal solution (I, ι) and some $a \in \mathbb{A}$.

Theorem 3.6 (Discretely self-similar spaces) *The following conditions on a topological space S are equivalent:*

- a. S is discretely self-similar
- b. S is the limit of some sequence $(\dots \longrightarrow S_2 \longrightarrow S_1)$ of finite discrete spaces
- c. S is the limit of some countable diagram of finite discrete spaces
- d. S is compact, metrizable, and totally disconnected.

Proof

(a) $\Rightarrow$ (b) Let $(\mathbb{A}, M)$ be a discrete self-similarity system. The universal solution is $\text{ob } \mathcal{I}$, and each $\text{ob } (\mathcal{I}a)$ is by definition the limit of a sequence of finite discrete spaces.

(b) $\Rightarrow$ (c) Trivial.

(c) $\Rightarrow$ (d) Compact metrizable spaces are the same as compact Hausdorff second countable spaces. The classes of compact Hausdorff spaces and totally disconnected spaces are closed under all limits, and the class of second countable spaces is closed under countable limits.

(d) $\Rightarrow$ (a) S has a basis $(U_n)_{n \geq 1}$ of open sets for which each U_n is also closed [Joh, Theorem II.4.2]. The separating sequence $(\mathcal{V}_n)_{n \geq 1}$ constructed in Lemma 3.2 then has the property that each $\mathcal{V}_n$ is a partition of S , so the resulting category $\mathbb{A}$ is discrete. This proves the implication when S is nonempty, and the empty case is trivial. $\square$

Example 3.7 The underlying topological space of the absolute Galois group $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ is a countable limit of finite discrete spaces, so discretely self-similar.

Define **finite self-similarity** as discrete self-similarity was defined, *mutatis mutandis*. Since there are uncountably many homeomorphism classes of compact metrizable spaces, most self-similar spaces are not finitely self-similar.

Conjecture 3.8 *The Julia set $J(f)$ of any complex rational function f is finitely self-similar.*

This brings us full circle: it says that in the first example of [Lei1], we could have taken any rational function f and seen the same type of behaviour: after a finite number of decompositions, no more new spaces I_n appear. Both $J(f)$ and its complement are invariant under f , so f restricts to an endomorphism of $J(f)$ and this endomorphism is, with finitely many exceptions, a $\deg(f)$ -to-one mapping. This suggests that f itself should provide the self-similarity structure of $J(f)$, and that if $(\mathbb{A}, M)$ is the corresponding self-similarity system then the sizes of $\mathbb{A}$ and M should be bounded in terms of $\deg(f)$.

4 The Cantor set

Three closely related classical results ([Wil], [HY]) describe the role of the Cantor set among compact metrizable spaces: up to homeomorphism,

- every totally disconnected compact metrizable space is a subspace of the Cantor set
- every nonempty compact metrizable space is a quotient of the Cantor set
- the only nonempty perfect totally disconnected compact metrizable space is the Cantor set.

These are all consequences of results on self-similarity systems.

We begin by studying discrete self-similarity systems, which, described as systems of equations [Lei1, eq. (6)], look something like this:

$$\begin{aligned} A &= A + A + B + D \\ B &= A + C + D \\ C &= C \\ D &= D + E \\ \vdots &\quad \vdots \end{aligned}$$




Figure 4-C: Definition of M'

By adjoining new objects, we can turn the system into one in which each right-hand side is a sum of at most two terms. Here, for instance, we can adjoin objects $A', A'', B', \dots$ and work with the system

$$A = A' + D, \quad A' = A'' + B, \quad A'' = A + A, \quad B = B' + D, \quad B' = A + C, \quad \dots$$

Lemma 4.1 (≤ 2 summands) *Let $(\mathbb{A}, M)$ be a discrete self-similarity system with universal solution (I, ι) . Then there is a discrete self-similarity system $(\mathbb{A}', M')$ with universal solution (I', ι') and an injection $\mathbb{A} \hookrightarrow \mathbb{A}'$ such that $I'|_{\mathbb{A}} \cong I$ and $|\sum_{b'} M'(b', a')| \leq 2$ for each $a' \in \mathbb{A}'$.*

Proof Given $a \in \mathbb{A}$, let $k(a) = |\sum_b M(b, a)|$ and write

$$\sum_b M(b, a) = \{b_1^a \xrightarrow{m_1^a} a, \dots, b_{k(a)}^a \xrightarrow{m_{k(a)}^a} a\}.$$

Let $\mathbb{A}' = \{(a, i) \mid a \in \mathbb{A}, 0 \leq i \leq k(a)\}$ and define $\mathbb{A} \hookrightarrow \mathbb{A}'$ by $a \mapsto (a, k(a))$. The module M' on $\mathbb{A}'$ has the property that $|M'(b', a')| \leq 1$ for each $b', a' \in \mathbb{A}'$, and is defined (Figure 4-C) by

$$|M'((b, j), (a, i))| = 1 \iff i \geq 1 \text{ and } (b, j) \in \{(a, i-1), (b_i^a, k(b_i^a))\}$$

$(b, a \in \mathbb{A}, 0 \leq j \leq k(b), 0 \leq i \leq k(a))$. Note that $(a, i-1) \neq (b_i^a, k(b_i^a))$.

Let (I', ι') be the universal solution of $(\mathbb{A}', M')$ in **Top**. I claim that for all $a \in \mathbb{A}$ and $i \in \{0, \dots, k(a)\}$,

$$I'(a, i) \cong I'(b_1^a, k(b_1^a)) + \dots + I'(b_i^a, k(b_i^a)).$$

This holds when $i = 0$ because $I'(a, 0) \cong (M' \otimes I')(a, 0) = \emptyset$. When $i \geq 1$,

$$I'(a, i) \cong (M \otimes I')(a, i) = I'(a, i-1) + I'(b_i^a, k(b_i^a)),$$

so the claim follows by induction. Taking $i = k(a)$, we have in particular $(I'|_{\mathbb{A}})a \cong (M \otimes (I'|_{\mathbb{A}}))a$ for each $a \in \mathbb{A}$.

This argument constructs an isomorphism $\psi : M \otimes (I'|_{\mathbb{A}}) \xrightarrow{\sim} I'|_{\mathbb{A}}$, described explicitly as follows. Any element of the module M is uniquely of the form

$b_i^a \xrightarrow{m_i^a} a$, and gives rise to a diagram

$$(b_i^a, k(b_i^a)) \dashrightarrow (a, i) \dashrightarrow \cdots \dashrightarrow (a, k(a)) \quad (3)$$

in $(\mathbb{A}', M')$, hence a diagram

$$I'(b_i^a, k(b_i^a)) \longrightarrow I'(a, i) \longrightarrow \cdots \longrightarrow I'(a, k(a)) \quad (4)$$

in **Top** in which the maps are components of $(\iota')^{-1}$; its composite is $\psi_{m_i^a} : (I'|_{\mathbb{A}})b_i^a \longrightarrow (I'|_{\mathbb{A}})a$.

We use the Precise Recognition Theorem to show that $(I'|_{\mathbb{A}}, \psi^{-1})$ is the universal solution of $(\mathbb{A}, M)$. It will follow that $I'|_{\mathbb{A}} \cong I$.

First, $I'|_{\mathbb{A}} : \mathbb{A} \longrightarrow \mathbf{Top}$ is a nondegenerate functor sending each $a \in \mathbb{A}$ to a compact space.

Second, $I'|_{\mathbb{A}}$ is occupied. For take a diagram $\cdots \xrightarrow{m_1} a_0$ in $(\mathbb{A}, M)$. Then as in (3), each m_i gives rise to a diagram

$$(a_{i+1}, k(a_{i+1})) \dashrightarrow \cdots \dashrightarrow (a_i, k(a_i))$$

in $(\mathbb{A}', M')$ of length at least 1. So there is an infinite diagram $\cdots \dashrightarrow (a_0, k(a_0))$ in $(\mathbb{A}', M')$. But I' is occupied, so $\emptyset \neq I'(a_0, k(a_0)) = (I'|_{\mathbb{A}})a_0$.

Third, $|\bigcap_{n \in \mathbb{N}} \psi_{m_1} \cdots \psi_{m_n}((I'|_{\mathbb{A}})a_n)| \leq 1$ for any diagram $\cdots \xrightarrow{m_1} a_0$ in $(\mathbb{A}, M)$. Indeed, each ψ_{m_i} is the composite (4) of one or more components of $(\iota')^{-1}$, so this follows from the corresponding property of $(\iota')^{-1}$. $\square$

A self-similarity system may contain equations of the form $A = 0$, that is, there may be objects $a \in \mathbb{A}$ for which there are no module elements $\cdot \dashrightarrow a$. If we are only interested in the self-similarity of nonempty spaces then we can discard those parts of the system, as follows.

Given a self-similarity system $(\mathbb{A}, M)$ (not necessarily discrete), write $\mathbb{A}^+$ for the full subcategory of $\mathbb{A}$ whose objects are those $a \in \mathbb{A}$ for which $\mathcal{I}a$ is nonempty, and M^+ for the restriction of M to $\mathbb{A}^+$.

Lemma 4.2 (≥ 1 summand) *Let $(\mathbb{A}, M)$ be a self-similarity system. Then $(\mathbb{A}^+, M^+)$ is a self-similarity system. Moreover, if $(\mathbb{A}, M)$ has a universal solution then so does $(\mathbb{A}^+, M^+)$, namely, the restriction of that of $(\mathbb{A}, M)$.*

Proof Finiteness of M^+ is trivial, and nondegeneracy follows from the isomorphism $\mathbb{E}(M^+(b, -)) \cong \mathbb{E}(M(b, -))$. So $(\mathbb{A}^+, M^+)$ is a self-similarity system.

Let (I, ι) be the universal solution of $(\mathbb{A}, M)$. ‘Moreover’ states that the restriction (I^+, ι^+) of (I, ι) to $(\mathbb{A}^+, M^+)$ is the universal solution of $(\mathbb{A}^+, M^+)$. For this restriction to make sense we need an isomorphism $M^+ \otimes (I|_{\mathbb{A}^+}) \cong (M \otimes I)|_{\mathbb{A}^+}$; indeed, if $a \in \mathbb{A}^+$ then

$$(M^+ \otimes (I|_{\mathbb{A}^+}))a = \int^{b \in \mathbb{A}^+} M^+(b, a) \times Ib \cong \int^{b \in \mathbb{A}} M(b, a) \times Ib = (M \otimes I)|_{\mathbb{A}^+}a.$$

To prove ‘moreover’, just observe that condition (c) (or (b)) of the Precise Recognition Theorem is satisfied by (I, ι) , hence by (I^+, ι^+) . $\square$

Corollary 4.3 (1 or 2 summands) *Let S be a nonempty discretely self-similar space. Then there is a discrete self-similarity system $(\mathbb{A}, M)$ with universal solution (I, ι) and an object $a \in \mathbb{A}$ such that $S \cong Ia$ and for each $a' \in \mathbb{A}$, $\sum_b M(b, a')$ has cardinality 1 or 2.*

Proof Take a self-similarity system from which S arises and apply Lemma 4.1 then Lemma 4.2. $\square$

Theorem 4.4 (Retracts of Cantor set) *Every nonempty discretely self-similar space is a retract of the Cantor set.*

Proof Let $(\mathbb{A}, M)$ be a discrete self-similarity system such that for each $a \in \mathbb{A}$, $\sum_b M(b, a)$ has cardinality 1 or 2, and choose for each $a \in \mathbb{A}$ a commutative triangle

$$\begin{array}{ccc} & 2 & \\ s_a \nearrow & & \searrow r_a \\ \sum_b M(b, a) & \xrightarrow{1} & \sum_b M(b, a) \end{array}$$

in **Set**. Write (I, ι) for the universal solution of $(\mathbb{A}, M)$. By the last corollary, it is enough to prove that each Ia is a retract of the Cantor set.

The Cantor set $2^{\mathbb{N}}$, together with the usual isomorphism $\gamma : 2^{\mathbb{N}} \xrightarrow{\sim} 2 \times 2^{\mathbb{N}}$, is the universal solution of the self-similarity system $(\mathbf{1}, 2)$ (Example 2.3). It gives rise to a functor $\Delta 2^{\mathbb{N}} : \mathbb{A} \longrightarrow \mathbf{Top}$ constant at $2^{\mathbb{N}}$ and a map

$$\varepsilon_a = \left(2^{\mathbb{N}} \xrightarrow[\sim]{\gamma} 2 \times 2^{\mathbb{N}} \xrightarrow{r_a \times 1} \sum_b M(b, a) \times 2^{\mathbb{N}} = (M \otimes \Delta 2^{\mathbb{N}})a \right)$$

for each $a \in \mathbb{A}$. So $(\Delta 2^{\mathbb{N}}, \varepsilon)$ is an M -coalgebra in **Top**, and there is a unique map $\rho : (\Delta 2^{\mathbb{N}}, \varepsilon) \longrightarrow (I, \iota)$ of M -coalgebras.

On the other hand, there is a map $\delta : \sum_a Ia \longrightarrow 2 \times \sum_a Ia$ whose a -component is the composite

$$Ia \xrightarrow[\sim]{\iota_a} \sum_b M(b, a) \times Ib \hookrightarrow \left(\sum_b M(b, a) \right) \times \left(\sum_b Ib \right) \xrightarrow{s_a \times 1} 2 \times \sum_b Ib.$$

There is a unique map $\sigma : (\sum_a Ia, \delta) \longrightarrow (2^{\mathbb{N}}, \gamma)$ of 2-coalgebras; preservation of the coalgebra structure says that for each $a \in \mathbb{A}$,

$$\begin{array}{ccc} Ia & \xrightarrow{\delta_a} & 2 \times \sum_b Ib \\ \sigma_a \downarrow & & \downarrow 1 \times \sigma \\ 2^{\mathbb{N}} & \xrightarrow{\gamma} & 2 \times 2^{\mathbb{N}} \end{array} \quad (5)$$

commutes. It follows that $(\sigma_a)_{a \in \mathbb{A}}$ is a map $(I, \iota) \longrightarrow (\Delta 2^{\mathbb{N}}, \varepsilon)$ of M -coalgebras, since for each $a \in \mathbb{A}$ there is a commutative diagram as in Figure 4-D.




Figure 4-D: Diagram for the proof of Theorem 4.4

We now have maps $(I, \iota) \xrightleftharpoons[\rho]{\sigma} (\Delta 2^{\mathbb{N}}, \varepsilon)$ of M -coalgebras, so by terminality of (I, ι) , the triangle

$$\begin{array}{ccc}
 & 2^{\mathbb{N}} & \\
 \sigma_a \nearrow & & \searrow \rho_a \\
 Ia & \xrightarrow{1} & Ia
 \end{array}$$

commutes for each $a \in \mathbb{A}$. □

Using Theorem 3.6, the first classical result follows:

Corollary 4.5 *Every nonempty totally disconnected compact metrizable space is a retract of the Cantor set.* □

We will deduce that every nonempty compact metrizable space is a quotient of the Cantor set. This is a topological analogue of the fact that colimits can be constructed from coproducts and coequalizers, or more precisely that every finite nonempty colimit is a coequalizer of maps between finite nonempty coproducts [Mac, V.2]. Recall from Example 2.3 that the Cantor set is isomorphic to the coproduct of k copies of itself for any $k \geq 1$, and is universal as such for any $k \geq 2$. Every compact metrizable space is self-similar, that is, defined by a system of finite colimit diagrams; if we decompose each colimit as a coequalizer

of coproducts, the coproduct part gives the Cantor set and the coequalizer gives the quotient.

Lemma 4.6 *Every self-similar space is a quotient of a discretely self-similar space.*

Proof Let $(\mathbb{A}, M)$ be a self-similarity system with universal solution (I, ι) , let $\mathbb{A}_0$ be the set of objects of $\mathbb{A}$ (regarded as a discrete category), and let M_0 be the restriction of M to $\mathbb{A}_0$. Then the universal solution of $(\mathbb{A}_0, M_0)$ is $(\text{ob } \mathbb{J}, \iota)$, and [Lei1, 6.7] says that Ia is a quotient of $(\text{ob } \mathbb{J})a$ for each $a \in \mathbb{A}$. $\square$

Theorem 4.7 (Quotients of Cantor set) *Every nonempty self-similar space is a quotient of the Cantor set.* $\square$

Corollary 4.8 *Every nonempty compact metrizable space is a quotient of the Cantor set.* $\square$

Finally, we deduce the characterization of the Cantor set as the unique nonempty perfect totally disconnected compact metrizable space. (A space is **perfect** if it is not homeomorphic to $T + \{\star\}$ for any space T , or equivalently if it has no isolated points.)

A discrete self-similarity system may contain equations of the form $A = A$, or loops such as $A = B, B = C, C = A$, or infinite chains such as $A_1 = A_2, A_2 = A_3, \dots$. In those cases the universal solution will involve the one-point space and perhaps other non-perfect spaces too (Example 2.4). But if the one-point space is not involved then the universal solution is extremely simple:

Proposition 4.9 (Empty or Cantor) *Let $(\mathbb{A}, M)$ be a discrete self-similarity system such that for all $a \in \mathbb{A}$, $|Ia| \neq 1$. Then each Ia is either empty or the Cantor set.*

Proof In Example 2.3 we saw that for $k \geq 2$, the universal solution of the self-similarity system $(\mathbf{1}, k)$ is $(2^{\mathbb{N}}, (\psi^{(k)})^{-1})$. Here $\psi_p^{(k)} : 2^{\mathbb{N}} \longrightarrow 2^{\mathbb{N}}$ is a contraction with constant $1/3$ for each p . Write $\psi_0^{(1)}$ for the identity on $2^{\mathbb{N}}$.

Define $J : \mathbb{A} \longrightarrow \mathbf{Top}$ by

$$Ja = \begin{cases} \emptyset & \text{if } Ia = \emptyset \\ 2^{\mathbb{N}} & \text{if } Ia \neq \emptyset. \end{cases}$$

For each $a \in \mathbb{A}$, let $k(a) = \left| \sum_{b: Ib \neq \emptyset} M(b, a) \right|$ and choose an isomorphism between the sets $\sum_{b: Ib \neq \emptyset} M(b, a)$ and $k(a)$; then for all $a \in \mathbb{A}$,

$$(M \otimes J)a = \sum_{b \in \mathbb{A}} M(b, a) \times Jb \cong \sum_{b: Ib \neq \emptyset} M(b, a) \times 2^{\mathbb{N}} \cong k(a) \times 2^{\mathbb{N}}.$$

Define an isomorphism $\gamma_a : Ja \xrightarrow{\sim} (M \otimes J)a$ for each $a \in \mathbb{A}$ as follows. If $Ia = \emptyset$ then $Ja = \emptyset = (M \otimes J)a$, so γ_a is determined. If $Ia \neq \emptyset$ then $Ja = 2^{\mathbb{N}}$ and we put

$$\gamma_a = \left(2^{\mathbb{N}} \xrightarrow[\sim]{(\psi^{(k(a))})^{-1}} k(a) \times 2^{\mathbb{N}} \cong (M \otimes J)a \right).$$

Certainly (J, γ) is a fixed point of M , J is occupied, and each Ja is compact and can be equipped with the usual metric. Now take a diagram

$$\cdots \xrightarrow{m_2} a_1 \xrightarrow{m_1} a_0.$$

For each $r \in \mathbb{N}$, either $k(a_r) = 1$, in which case $(\gamma^{-1})_{m_{r+1}}$ is an isometry, or $k(a_r) \geq 2$, in which case $(\gamma^{-1})_{m_{r+1}}$ is a contraction with constant $1/3$. The latter case arises infinitely often: for if not, there is some $s \in \mathbb{N}$ for which $1 = k(a_s) = k(a_{s+1}) = \cdots$, and then $|Ia_s| = 1$, contrary to hypothesis. Moreover, $\text{diam}(Ja_r) \leq 1$ for each r . So condition (c) of the Precise Recognition Theorem is satisfied and (J, γ) is the universal solution. $\square$

We also need an intuitively obvious lemma: that to calculate the value of the universal solution at a , we can ignore all parts of the system that do not contribute to a . Precisely, let $(\mathbb{A}, M)$ be a self-similarity system and $a \in \mathbb{A}$. Write $\mathbb{A}^{(a)}$ for the full subcategory of $\mathbb{A}$ whose objects are those $b \in \mathbb{A}$ for which there exists a diagram

$$b = a_n \xrightarrow{m_n} \cdots \xrightarrow{m_1} a_0 = a \quad (6)$$

for some $n \in \mathbb{N}$, and $M^{(a)}$ for the restriction of M to $\mathbb{A}^{(a)}$.

Lemma 4.10 *Let $(\mathbb{A}, M)$ be a self-similarity system and $a \in \mathbb{A}$. Then $(\mathbb{A}^{(a)}, M^{(a)})$ is a self-similarity system. Moreover, if $(\mathbb{A}, M)$ has a universal solution then so does $(\mathbb{A}^{(a)}, M^{(a)})$, namely, the restriction of that of $(\mathbb{A}, M)$.*

Proof Finiteness of $M^{(a)}$ is trivial. For nondegeneracy, observe that if $b' \longrightarrow b$ in $\mathbb{A}$ and $b \in \mathbb{A}^{(a)}$ then $b' \in \mathbb{A}^{(a)}$; it follows that $M^{(a)}$ satisfies **ND1** and **ND2**. So $(\mathbb{A}^{(a)}, M^{(a)})$ is a self-similarity system.

‘Moreover’ states that if (I, ι) is the universal solution of $(\mathbb{A}, M)$ then $(I^{(a)}, \iota^{(a)})$ is the universal solution of $(\mathbb{A}^{(a)}, M^{(a)})$, where $I^{(a)}$ is the restriction of I and $\iota^{(a)}$ is the restriction of ι . For the latter to make sense we need an isomorphism $(M \otimes I)|_{\mathbb{A}^{(a)}} \cong M \otimes (I|_{\mathbb{A}^{(a)}})$; indeed, the square

$$\begin{array}{ccc} [\mathbb{A}, \mathbf{Set}] & \xrightarrow{M \otimes -} & [\mathbb{A}, \mathbf{Set}] \\ \text{restriction} \downarrow & & \downarrow \text{restriction} \\ [\mathbb{A}^{(a)}, \mathbf{Set}] & \xrightarrow{M^{(a)} \otimes -} & [\mathbb{A}^{(a)}, \mathbf{Set}] \end{array}$$

commutes up to canonical isomorphism, as if $X \in [\mathbb{A}, \mathbf{Set}]$ and $a' \in \mathbb{A}^{(a)}$ then

$$\begin{aligned} (M^{(a)} \otimes (X|_{\mathbb{A}^{(a)}})) a' &= \int^{b \in \mathbb{A}^{(a)}} M^{(a)}(b, a') \times Xb \\ &\cong \int^{b \in \mathbb{A}} M(b, a') \times Xb \\ &= (M \otimes X) a'. \end{aligned}$$

To prove ‘moreover’, just observe that condition (c) (or (b)) of the Precise Recognition Theorem is satisfied by (I, ι) , hence by $(I^{(a)}, \iota^{(a)})$. $\square$

Theorem 4.11 (Characterization of Cantor set) *A perfect discretely self-similar space is either empty or the Cantor set.*

Proof By the previous lemma we may assume that the space is of the form Ia where $(\mathbb{A}, M)$ is a discrete self-similarity system with universal solution (I, ι) , $a \in \mathbb{A}$, and for all $b \in \mathbb{A}$ there exists a finite chain (6). But

$$Ia \cong \sum_{a_1} M(a_1, a) \times Ia_1 \cong \sum_{a_1, a_2} M(a_1, a) \times M(a_2, a_1) \times Ia_2 \cong \cdots,$$

so for each $b \in \mathbb{A}$, Ib is a summand of Ia ; and since Ia is perfect, $|Ib| \neq 1$. The result follows from Proposition 4.9. $\square$

Corollary 4.12 *A perfect totally disconnected compact metrizable space is either empty or the Cantor set.* $\square$

References

- [Bou1] Nicolas Bourbaki, *General Topology* Part 1, Springer, 1966.
- [Bou2] Nicolas Bourbaki, *General Topology* Part 2, Springer, 1966.
- [Freu] Hans Freudenthal, Simplicialzerlegungen von beschränkter Flachheit, *Annals of Mathematics* (2) 43 (1942), 580–582.
- [Hat] Allen Hatcher, *Algebraic Topology*, Cambridge University Press, 2002; also available from www.math.cornell.edu/~hatcher.
- [HY] John G. Hocking, Gail S. Young, *Topology*, Addison–Wesley, 1961.
- [Hut] John E. Hutchinson, Fractals and self-similarity, *Indiana University Mathematics Journal* 30 (1981), no. 5, 713–747.
- [Joh] Peter T. Johnstone, *Stone Spaces*, Cambridge Studies in Advanced Mathematics 3, Cambridge University Press, 1982.
- [Lei1] Tom Leinster, A general theory of self-similarity I, e-print math.DS/0411344, 2004.
- [Lei2] Tom Leinster, General self-similarity: an overview, e-print math.DS/0411343, 2004.
- [Mac] Saunders Mac Lane, *Categories for the Working Mathematician*, Graduate Texts in Mathematics 5, Springer, 1971.
- [Wil] Stephen Willard, *General Topology*, Addison–Wesley, 1970.